

Upper Endoscopy Prep Instructions

7 days prior to procedure:

- Please confirm a ride home from your procedure, as you will receive medicine for sedation.
 - Hospital policy requires that you have a ride home after your procedure by a responsible adult. Taxis, Uber's, Lyft's or public transportation are not recommended.

Failure to obtain a ride home may lead to cancellation of your procedure.

1 day prior to procedure:

• No solid food after midnight. Drink only clear liquids from the guidelines below.

DO NOT consume:

- Liquids that are **red**, orange, or purple
- o Alcohol
- Dairy or dairy substitutes
- Solid food
- Juices you cannot see through
- o Soup
- Protein drinks



- o Sports drinks
- Clear juice, such as apple or white grape
- o Clear carbonated beverages such as seltzer, ginger ale, Sprite or cola
- o Water
- Black coffee or tea
- Popsicles

Day of Procedure:

- 3 hours before procedure arrival time:
 - **STOP** drinking all liquids, including the bowel prep.
 - No chewing gum or hard candy.
 - \circ You may take your morning medications with a sip of water at this time.
 - You may take your morning medications with a sip of water at this time.



Additional Information

Ride Information

- You can expect to be ready to go home 1¹/₂ to 2¹/₂ hours after your arrival.
- After being dropped off, please be sure your ride is reachable by phone and able to return to the hospital within 30 minutes of receiving a call from us.
- Cell phone reception is limited in the parking garage. If your ride plans to stay in the car, it is best to park on the top floor of the garage.
- When your ride arrives to pick you up, please have them pull up to the hospital main entrance and provide your name to the valet. You will be transported via wheelchair to your ride.

Important Information on Medications

If you are taking a blood thinner, (examples; Coumadin, Plavix, Xarelto), you will receive instructions from our scheduling team about when to stop this medication, ordered by your prescribing provider.

If you are taking a diabetes medication, please call your prescribing provider for instructions before your procedure.

**Glucagon-like peptide agonists (GLP-1) is group of medication used for diabetes or weight loss (examples; Ozempic, Wegovy, Saxenda, Byetta). If you are taking a GLP-1 medication, please call your prescribing provider about when to stop them as they can slow stomach emptying.

Failure to follow medication instructions may lead to cancellation of your procedure.

• If you wear contacts, you will be asked to remove them before the procedure.



- If you are within childbearing age, you may be asked for a urine sample before to your procedure.
- Please ignore the procedure times listed on Exeter Hospital or Core Physicians' patient portals. The Endoscopy department schedulers will contact you directly with your time.
- You will be asked to remove jewelry before your procedure.
- You should consider leaving valuable items at home.

Please scan the QR code with your cell phone to view the Endoscopy Department's Prep Instruction Video for further information.

For questions, or if you are unable to attend your appointment, please call the endoscopy department at 603-580-7892.